


CB660SP



DESCRIPTION

NEURAL STEM CELLS DERIVED FROM HUMAN FOETAL SPINAL CORD

Organism:	<i>Homo sapiens</i> , human
Cell Type:	Human neural stem cell
Source:	Foetal spinal cord at embryonic 50-55 days (Carnegie stage 19-22)
Datasheet:	available under request

REFERENCES

1. Sun Y, Pollard S, Conti L, Toselli M, Biella G, Parkin G, Willatt L, Falk A, Cattaneo E, Smith A. Long-term tripotent differentiation capacity of human neural stem (NS) cells in adherent culture. *Mol Cell Neurosci*. 2008 Jun;38(2):245-58.
2. Sun Y, Kong W, Falk A, Hu J, Zhou L, Pollard S, Smith A. CD133 (Prominin) negative human neural stem cells are clonogenic and tripotent. *PLoS One*. 2009;4(5):e5498.
3. Fujimoto Y, Abematsu M, Falk A, Tsujimura K, Sanosaka T, Juliandi B, Semi K, Namihira M, Komiya S, Smith A, Nakashima K. Treatment of a mouse model of spinal cord injury by transplantation of human induced pluripotent stem cell-derived long-term self-renewing neuroepithelial-like stem cells. *Stem Cells*. 2012 Jun;30(6):1163-73.
4. Baronchelli S, Bentivegna A, Redaelli S, Riva G, Butta V, Paoletta L, Isimbaldi G, Miozzo M, Tabano S, Daga A, Marubbi D, Cattaneo M, Biunno I, Dalprà L. Delineating the cytogenomic and epigenomic landscapes of glioma stem cell lines. *PLoS One* 2013;8(2):e57462.